



Volatility modeling in financial markets

Master Thesis

Sergiy Ladokhin

Supervisors:

Dr. Sandjai Bhulai, VU University Amsterdam

Brian Doelkahar, Fortis Bank Nederland

VU University Amsterdam

Faculty of Sciences, Business Mathematics and Informatics

De Boelelaan 1081a, 1081 HV Amsterdam

Host organization:

Fortis Bank Nederland, Brokerage Clearing and Custody

Rokin 55, 1012 KK Amsterdam

PUBLIC VERSION

26 July 2009

Summary

This project focuses on the problem of volatility modeling in financial markets. It begins with a general description of volatility and its properties, and discusses its usage in financial risk management. The research is divided into two parts: estimation of conditional volatility and modeling of volatility skews. The first one is focused on comparing different models for conditional volatility estimation. We examine the accuracy of several of the most popular methods: historical volatility models (e.g., Exponential Weighted Moving Average), the implied volatility, and autoregressive conditional heteroskedastic models (e.g., the GARCH family of models). The second part of the project is dedicated to modeling the implied volatility skews and surfaces. We introduce a number of representations of the volatility skews and discuss their importance for the risk management of the options portfolio. The comparison analysis of several approaches to the volatility skews modeling (including spline models and the SABR family of models) is made. Special attention is paid to modeling the dynamics of the implied volatility surfaces in time.

This research is done for the Fortis Bank Nederland Brokerage, Clearing and Custody (FBNBCC). All of the models and methods described in this research are designed to improve the methodology currently used by FBNBCC. The models of this study were implemented, calibrated, and tested using real market data; and their results were compared to the currently used methods by the FBNBCC's risk management system. Another objective of this study is to examine potential shortcomings of FBNCC's risk management system and to develop recommendations for their elimination.

Table of Contents

Summary	2
1 Introduction.....	4
2 Comparison analysis of models for volatility forecasting	6
2.1 Role of volatility in the estimation of the market risk	6
2.2 Metrics for the market risk	8
2.3 Models for conditional volatility	11
2.4 Applications of Extreme Value Theory to the market risk	18
2.5 Comparison of the volatility forecasting models	20
3 Modeling implied volatility surfaces	24
3.1 Risk of volatility “smile” and its impact.....	24
3.2 Representation of the moneyness.....	25
3.3 Models of the implied volatility surface.....	27
3.4 Modeling the dynamics of volatility surfaces.....	35
3.5 Comparison of the volatility surface models	40
4 Conclusions and practical recommendations	45
5 Appendixes	46
5.1 t Location-Scale distribution	46
5.2 The Black-Scholes option pricing equation	48
5.3 Cross- validation	50
5.4 Results of the conditional volatility estimation	51
5.5 Results of implied volatility surface modeling	55
6 Bibliography	58

1 Introduction

The main characteristic of any financial asset is its return which is typically considered to be a random variable. The spread of outcomes of this variable, known as assets volatility, plays an important role in numerous financial applications. Its primary usage is to estimate the value of market risk. Volatility is also a key parameter for pricing financial derivatives. All modern option-pricing techniques rely on a volatility parameter for price evaluation. Volatility is also used for risk management assessment and in general portfolio management. It is crucial for financial institutions not only to know the current value of the volatility of the managed assets, but also to be able to predict their future values. Volatility forecasting is especially important for institutions involved in options trading and portfolio management.

Accurate estimation of the future behavior of the values of financial indicators is obscured by complex interconnections between these indicators, which are often convoluted and not intuitive. This makes forecasting the behavior of volatility a challenging task even for experts in this field. Mathematical modeling can assist in establishing the relationship between current values of the financial indicators and their future expected values. Model-based quantitative forecasts can provide financial institutions with a valuable estimate of a future market trend. Although some experts believe that future events are unpredictable, some empirical evidence to the contrary exists. For example, financial volatility has a tendency to cluster and exhibits considerable autocorrelation (i.e., the dependency of future values on past values). These features provide the justification for formalizing the concept of volatility and creating mathematical techniques for volatility forecasting. Starting from the late 70's a number of models for volatility forecasting have been introduced.

The purpose of this project is to compare different mathematical methods used in modeling the volatility of different assets. The thesis is divided into two parts: comparison of the volatility estimation methods and modeling volatility skews. The first one is focused on introducing the general framework of dynamic risk management and comparison of different models for volatility forecasting. Specifically, we tested several classes of volatility forecasting models that are widely used in modern practice: historical (including moving averages), autoregressive, conditional heteroscedastic models, and the implied volatility concept. Moreover, we introduced a model "blending" procedure which can potentially improve individual "classic" methods. The second part of this work is dedicated to modeling the implied volatility surfaces. This problem plays a key role in

managing the risk of options portfolios. We discussed and compared several models of the approximation of the surfaces, as well as several approaches to the dynamics of these surfaces. All of the models and algorithms discussed in this work are tested using different classes of market data.

2 Comparison analysis of models for volatility forecasting

2.1 Role of volatility in the estimation of the market risk

Market risk is one of the main sources of uncertainty for any financial institution that holds risky assets. In general, market risk refers to the possibility that the portfolio value will decrease due to the changes in market factors. An example of market factors is a change of the price of securities, indices of securities, changes in interest rates, currency rates, etc. The market risk has a significant influence on the value of the exposed financial institution. Unpredicted changes in the market situation can potentially lead to big losses; therefore the market risk must be estimated by any institution involved in security markets.

There are a number of approaches to estimate the exposure of the financial institution to the market risk. The Value-at-Risk methodology is the most heavily used one for the estimates of the market risk in practical applications. The concept of volatility plays a key role in this methodology. Volatility of the asset refers to the uncertainty of the value of the returns from holding risky assets over a given period of time. Correct estimation of the volatility can provide a substantial advantage to the financial institution. The parameters of the Value-at-Risk methodology could be estimated over different time periods (e.g., yearly, monthly, weekly, etc.). The estimate made on a daily basis is the most adequate, because the market situation changes very rapidly. Dynamic Risk management is the technique that monitors the market risk on the daily basis. Dynamic Risk Management requires not only the correct estimate of the historical volatility, but also a short term forecast. This forecast sometimes is referred as conditional volatility estimation. For the last 30 years a number of successful volatility models were developed.

The purpose of this chapter is to compare different methods for conditional volatility estimation (forecasting). This comparison will be made with the respect to the goals of the dynamic risk management. We will start this assessment with the introduction of a general framework of risk management, its metrics and methodology. Then we will discuss the comparison of the volatility models among themselves. In particular, we will test several classes of volatility forecasting models that are widely used in modern practice: historical (including exponential moving average), autoregressive, conditional heteroscedastic models, and the implied volatility concept. In addition, we examine a relatively new approach: a model blending technique. Its ability to overcome the

disadvantages of single models by combining them is a feature, which can give an improvement of volatility forecasting. All of the models were tested on daily data from different asset classes.

2.2 Metrics for the market risk

In this chapter we will introduce the general framework for measuring the market risk, but first we have to give a few basic definitions. We will be interested in the risk of holding some risky asset S , which price for day t is S_t . Let us assume that the price of the asset is positive $S_t > 0$. The return of holding such an asset is given by:

$$r_t = \ln\left(\frac{S_t}{S_{t-1}}\right). \quad (2.1)$$

The return is considered to be a random variable. We will be interested in a time series of the returns over some time period. The return is characterized by the expected value μ and volatility σ . The expected value of the return at each given time t could be taken as zero, making volatility σ the most important characteristic of the return. Volatility refers to the spread of all outcomes of an uncertain variable. In finance, we are interested in the outcomes of asset returns. Volatility is associated with the sample standard deviation of returns over some period of time. It is computed using the following equation:

$$\hat{\sigma} = \sqrt{\frac{1}{T-1} \sum_{t=1}^T (r_t - \mu)^2}, \quad (2.2)$$

where r_t is the return of an asset over period t and μ is the average return over T periods.

The variance, σ^2 , could also be used as a measure of volatility. But this is less common, because variance and standard deviation are connected by a simple relationship. Volatility is a quantified measure of market risk. Volatility is related to risk, but it is not exactly the same. Risk is the uncertainty of a negative outcome of some event (e.g., stock returns); volatility measures the spread of outcomes. This includes positive as well as negative outcomes.

Also, we will be interested in a loss function which describes the negative outcomes of the returns. Let us denote by V_t the value of some position (single stock, index, currency, etc.) on day t . The logarithmic return on the next day is r_{t+1} , so the loss over the next day is $L_{t+1} = l(r_{t+1}) = -V_t r_{t+1}$. For the sake of simplicity we can assume $V_t = 1$ for all t . Simply speaking, the loss function is a function of negative log-returns. The loss function is introduced accordingly to (McNeil F. &, 2005).

The Value-at-Risk (VaR) is probably the most widely used metrics to measure the market risk. Let us consider a portfolio of risky assets, and denote by $F_L(l) = P(L \leq l)$ the cumulative distribution function of the corresponding loss distribution. The VaR could be viewed as a maximum loss of the given portfolio which is not exceeded with a given high probability (McNeil F. &, 2005). Usually, the Value-at-Risk is computed for some confidence level $\alpha \in (0,1)$. The VaR of the portfolio at confidence level α is given by the smallest number l such that the probability that the loss L exceeds l is no larger than $(1 - \alpha)$:

$$VaR_\alpha = \inf\{l \in \mathbb{R}: P(L > l) \leq 1 - \alpha\} = \inf\{F_L(l) \geq \alpha\}. \quad (2.3)$$

From a statistical point of view, the VaR is a quantile of a loss distribution. The definition of VaR implies that we do not know anything about the size of losses that exceeds the given threshold. This is one of the major disadvantages of the VaR as a measure of risk. The Expected Shortfall (ES) was introduced to overcome these difficulties. For the loss L and cumulative distribution function F_L the expected shortfall at confidence level $\alpha \in (0,1)$ is defined as

$$ES_\alpha = \frac{1}{1 - \alpha} \int_\alpha^1 VaR_u(L) du. \quad (2.4)$$

Alternatively, we can define ES as a loss L that is realized in the event that the VaR is exceeded:

$$ES_\alpha = E(L|L \geq VaR_\alpha). \quad (2.5)$$

The proof of (2.5) can be found in (McNeil F. &, 2005).

Conditional risk management plays a key role for the purposes of financial clearing. By conditional risk management we will understand the re-computing of key risk measures (the VaR and the ES) on a daily basis, given the changes in the market situation. The conditional loss process for day t is modeled by the following equation:

$$L_t = \mu_t + \sigma_t Z_t, \quad (2.6)$$

where Z_t are the random residuals with expected value zero and variance 1. We will assume that the distribution of the residuals has a cumulative distribution function G . The general equation for the Value-at-Risk and the Expected Shortfall is:

$$VaR_\alpha^t = \mu_{t+1} + \sigma_{t+1} q_\alpha(Z), \quad (2.7)$$

$$ES_\alpha^t = \mu_{t+1} + \sigma_{t+1} ES_\alpha(Z), \quad (2.8)$$

where $q_\alpha(Z)$ is a quantile of the distribution of residuals, $ES_\alpha(Z)$ is the corresponding expected shortfall and $\alpha \in (0,1)$ is a given confidence level. We will require estimates of the conditional mean μ_{t+1} and conditional volatility σ_{t+1} in order to use the above equations. Moreover, the model of the distribution of the residuals Z has to be build to estimate the quantile and Expected Shortfall of Z .

In practical applications the conditional mean is usually taken to be equal to zero $\mu_{t+1} = 0$. It will simplify equations (2.7)-(2.8):

$$VaR_\alpha^t = \sigma_{t+1}q_\alpha(Z), \quad (2.9)$$

$$ES_\alpha^t = \sigma_{t+1}ES_\alpha(Z). \quad (2.10)$$

The estimate of the conditional variance σ_{t+1} can be done by different methods. An overview of some of these models will be given in the next section.

2.3 Models for conditional volatility

Estimating the conditional volatility σ_{t+1} is an important element of dynamic risk management. We can refer to this problem as volatility forecast, because we have to estimate the volatility at time $t + 1$ given data up to time t . Formally, forecasting the volatility could be seen as finding $\hat{\sigma}_t$ that will minimize the error $\varepsilon = f(\sigma_t - \hat{\sigma}_t)$, where σ_t is an actual (or observed) volatility over period t and $f(\cdot)$ is an error function. A discussion of different forms of error functions can be found below. To estimate the volatility on a certain timeframe one could use data of a smaller timeframe and compute the standard deviation. For example, if we are interested in the monthly volatility, we can compute the standard deviation of daily returns. Sometimes it is difficult to find data for a shorter timeframe, in which case different methods for volatility estimation can be used. The simplest and, perhaps the most effective, way of estimating the volatility is taking the daily squared returns as a proxy of the conditional variance $\sigma_t^2 \equiv r_t^2$.

In order to estimate the forecasting performance of some methods or to compare several methods we should define error functions. Although, the error function can be defined in a number of ways, we will focus on two of them the Root Mean Square Error (RMSE) and the Mean Heteroscedastic Square Error (MHSE). For more information about different error functions see, for example (Poon S. H., 2005). The Root Mean Square Error is given by:

$$RMSE = \sqrt{\frac{1}{N} \sum_{t=1}^N (\hat{\sigma}_t - \sigma_t)^2}, \quad (2.11)$$

where σ_t is the observed volatility (absolute value of returns) on a day t , $\hat{\sigma}_t$ is a forecast of the volatility and N is the number of days in the given data set. Similarly, the Mean Heteroscedastic Square Error is given by the following equation:

$$MHSE = \frac{1}{N} \sum_{t=1}^N \left(\frac{\sigma_t}{\hat{\sigma}_t} - 1 \right)^2. \quad (2.12)$$

The main difference between the RMSE and the MHSE is, that the RMSE measures the error in terms of average deviations and the HMSE as an average relative error. Volatility is not constant over time. Moreover it exhibits certain patterns. This means that large movements in returns tend to be followed by further large movements. Thus the economy has cycles with high volatility and low

volatility periods. The RMSE is a very popular error function among practitioners; however, it is not always the best one, especially when volatility clustering occurs. Obviously the RMSE is an important measure, but it is not always sufficient for accurate model comparison. For example, one forecasting method can systematically underestimate volatility, while the other will overestimate. The RMSE of these two methods will be the same, but clearly, the second method is more preferred to the first. The accuracy of the VaR and the ES (2.9)-(2.10) should be also taken into consideration in order to compare different forecasting methods.

In this section we will discuss several methods for volatility forecasting. We will intentionally skip the discussion of the simple models (e.g., Simple Moving Average) and focus on models that have a proven forecasting power, such as Exponentially Weighted Moving Average, Autoregressive Conditional Heteroskedasticity, Generalized Autoregressive Conditional Heteroskedasticity, and others. We will also introduce blending procedures which are aimed to overcome disadvantages of individual models.

We will start our discussion with the **Exponentially Weighted Moving Average (EWMA)**: an estimation method suggested by the Risk Metrics framework (J.P. Morgan/Reuters, 1996). The volatility forecast for day $t + 1$ is given by the following equation:

$$\hat{\sigma}_{t+1} = \sqrt{(1 - \lambda) \sum_{i=1}^n \lambda^{i-1} (r_{t+1-i} - \bar{r})^2}, \quad (2.13)$$

where

λ ($0 < \lambda < 1$) the parameter of the model, so called decay factor,

r_{t+1-i} , $i = 1, \dots, n$ previously observed returns,

\bar{r} is an exponentially weighted moving average mean of the daily returns, and it is given by the following equation

$$\bar{r} = (1 - \lambda) \sum_{i=1}^n \lambda^{i-1} r_{t+1-i}. \quad (2.14)$$

An attractive feature of the EWMA model is that it can be rewritten in recursive form. In order to do this, we have to assume that an infinite number of historic data is available. Then (2.13) can be rewritten as:

$$\hat{\sigma}_{t+1} = \sqrt{\lambda \hat{\sigma}_t^2 + (1 - \lambda)r_t^2}, \quad (2.15)$$

where $\hat{\sigma}_t^2$ is an EWMA estimate of the variance for the previous day. This representation is very efficient for the computational purposes.

There are a number of different methods for the calibration of the parameters of the EWMA model. An extensive overview of these approaches is given in (J.P. Morgan/Reuters, 1996). We will briefly discuss the main ideas of these approaches. The parameter n refers to the number of historical observations used to produce the estimate. A different estimate of this parameter does not significantly influence the accuracy of forecast. The Risk Metrics framework (J.P. Morgan/Reuters, 1996) suggests taking this parameter equal to 125. On the contrary, the λ parameter has a much more significant influence on the quality of forecast. The decay factor λ can be interpreted as a weight, which is given to the last observed volatility. Usually λ is taken to be close to 1. With the smaller values of λ the EWMA reacts more to the recent changes of the observed volatility, while with the bigger value the EWMA tends to “smooth” the observations more. The Risk Metrics suggests taking $\lambda = 0.94$. The optimal value of the decay factor, λ^* , can be found as a result of an optimization procedure. We will maximize the likelihood function to estimate λ . First, we have to assume some distribution of the returns. We will use the t-Location Scale Distribution as a distribution of the daily returns. The detailed description of the t-Location Scale Distribution and a rational behind the decision to use it is given in Appendix 5.1. For practical considerations we can minimize the negative log likelihood instead of maximizing the likelihood function. The procedure is then given by the following equation:

$$\lambda^* = \arg \min_{0 < \lambda < 1} \left[- \sum_{i=1}^N \log f_{\lambda}(r_i) \right], \quad (2.16)$$

where

$f_{\lambda}(\cdot)$ is a probability density function of a t-Location Scale distribution which depends on parameter λ ,

r_i are historical returns,

N is the number of observed returns in a given data set.

The optimization procedure (2.16) becomes easy to implement and apply, after the parameters of the distribution of returns are determined.

The financial market volatility is known to cluster (Tsay, 2005). A highly volatile period tends to persist for some time before the market returns to a more stable environment. An autoregressive approach helps to build more accurate and reliable volatility models.

The **Autoregressive Conditional Heteroskedasticity (ARCH)** model was first introduced by Engle in 1982 (Engle, 1982). The ARCH model and its extensions (GARCH, EGARCH, etc.) are among the most popular models for forecasting market returns and volatility. Originally, the ARCH model rather than using standard deviations used the variance. Let us call the variance of the returns σ^2 as h . The ARCH model can be defined as follows:

$$r_{t+1} = \mu + \varepsilon_{t+1}, \quad (2.17)$$

$$\varepsilon_{t+1} = \sqrt{h_{t+1}} z_{t+1}, \quad (2.18)$$

$$h_{t+1} = \alpha_0 + \sum_{j=1}^q \alpha_j \varepsilon_{t+1-j}^2, \quad (2.19)$$

where

r_{t+1} is the conditional estimate of returns at time $t + 1$,

μ is the mean return. As was mentioned earlier, it can be taken to be equal to zero,

ε_{t+1} are the residuals (or error terms),

$z_{t+1} \sim iid N(0,1)$ normally distributed random variables,

$\alpha_0, \alpha_1, \alpha_2, \dots, \alpha_q$ are parameters of the model.

We will refer to this process as an ARCH(q) process. The process z_{t+1} is scaled by h_{t+1} , the conditional variance, which follows an autoregressive regression process. The parameters $\alpha_j \geq 0, j = 1, \dots, q$ insure that the variance h_{t+1} is positive. The one step ahead forecast is simply the square root of the variance $\hat{\sigma}_{t+1} = \sqrt{h_{t+1}}$. The parameter q is usually taken to be 1 or 2. Higher orders of the ARCH model are less effective (Tsay, 2006). The name, ARCH, refers to this structure: the model is autoregressive, since ε_t clearly depends on previous ε_{t-i} , and conditionally heteroscedastic, since the conditional variance changes continually.

The estimates of the parameters $\alpha_0, \alpha_1, \alpha_2, \dots, \alpha_q$ are made in a similar way as in the case of the EWMA model. We will search for such values of parameters, which minimize the negative log-likelihood function. This procedure is similar to (2.16).

The **Generalized Autoregressive Conditional Heteroskedasticity (GARCH)** model is a general version of the ARCH model. It differs from ARCH by the form of h_{t+1} . Formally, the $GARCH(p,q)$ model can be defined as follows:

$$r_{t+1} = \mu + \varepsilon_{t+1}, \quad (2.20)$$

$$\varepsilon_{t+1} = \sqrt{h_{t+1}}z_{t+1}, \quad (2.21)$$

$$h_{t+1} = \omega + \sum_{i=1}^p \beta_i h_{t+1-i} + \sum_{j=1}^q \alpha_j \varepsilon_{t+1-j}^2, \quad (2.22)$$

where

r_{t+1} is the conditional estimate of returns at time $t + 1$,

μ is the mean return. Again, it can be taken to be equal to zero, $\mu = 0$,

ε_{t+1} are the residuals (or error terms),

$z_{t+1} \sim iid N(0,1)$ are normally distributed random variables,

$\omega, \alpha_1, \alpha_2, \dots, \alpha_q, \beta_1, \beta_2, \dots, \beta_p$ are parameters of the model.

As before, parameters $\omega \geq 0, \beta_i \geq 0, \alpha_j \geq 0$ are positive. There are additional constraints on β_i, α_j for models with higher orders than $GARCH(1,1)$ (Tsai, 2006). As in the ARCH model, at time t all the parameters are known, and h_{t+1} can be easily computed. The one-step ahead forecast of the volatility is again, just $\hat{\sigma}_{t+1} = \sqrt{h_{t+1}}$. The parameters $\omega, \alpha_1, \alpha_2, \dots, \alpha_q, \beta_1, \beta_2, \dots, \beta_p$ of the model can be found by algorithm (2.16).

There are a number of extensions of the GARCH model, such as Integrated GARCH, Exponential GARCH, GJR- GARCH and others [J. Knight 2007]. We will include in our analysis the **Exponential Generalized Autoregressive Conditional Heteroskedasticity (EGARCH)**, which is perhaps, the most widely used extension of GARCH in practical applications. The $EGARCH(p,q)$ model is given as follows:

$$r_{t+1} = \mu + \varepsilon_{t+1}, \quad (2.23)$$

$$\varepsilon_{t+1} = \sqrt{h_{t+1}}z_{t+1}, \quad (2.24)$$

$$\log h_{t+1}^2 = \omega + \sum_{i=1}^p \beta_i \log h_{t+1-i}^2 + \sum_{j=1}^q \alpha_j \frac{|\varepsilon_{t+1-j}|}{h_{t+1-j}}, \quad (2.25)$$

where

r_{t+1} is the conditional estimate of the returns at time $t + 1$,
 μ is the mean return. Again, it can be taken to be equal zero, $\mu = 0$,
 ε_{t+1} are the residuals (or error terms),
 $z_{t+1} \sim iid N(0,1)$ are normally distributed random variables,
 $\omega, \alpha_1, \alpha_2, \dots, \alpha_q, \beta_1, \beta_2, \dots, \beta_p$ are parameters of the model.

The EGARCH has similar properties as GARCH model. The conditional volatility estimate is, again given by $\hat{\sigma}_{t+1} = \sqrt{h_{t+1}}$. For a detailed description of these models see (Nelson, 1991).

Model Blending is a popular statistical technique to increase the forecasting power of models (Witten, 2005). It is applied when there are a number of models that estimate the same parameter. Some models tend to underestimate the real value of the forecasting parameter; in contrast others tend to overestimate. Model blending is an approach to overcome disadvantages of individual models and combine their advantages. We will consider the linear model blending. Formally it can be given as follows:

$$\hat{\sigma}_{t+1} = \alpha_0 + \alpha_1 P_1(d_t, \gamma_1) + \alpha_2 P_2(d_t, \gamma_2) + \dots + \alpha_k P_k(d_t, \gamma_k), \quad (2.26)$$

where

P_1, P_2, \dots, P_k are the individual models for the volatility forecast,
 $\gamma_1, \gamma_2, \dots, \gamma_k$ is the set of optimal parameters for each of the volatility models,
 d_t is necessary historical data as input of the volatility models,
 $\alpha_0, \alpha_1, \alpha_2, \dots, \alpha_k$ are parameters of the models.

We will use the following models as $P_i, i = 1, \dots, k$: *EWMA*, *ARCH*(q), *GARCH*(p, q), and *EGARCH*(p, q). The parameters $\alpha_0, \alpha_1, \alpha_2, \dots, \alpha_k$ of the model blending can be optimized in several ways. We will consider two ways of optimizing these parameters, both of them are to minimize the error function. The first one is to minimize the error function in the form of the RMSE (2.11) between the observed volatilities and the outputs of the model (2.26). The second one is determines estimates of the parameters by minimizing the MHSE (2.12). These two different approaches lead to two different sets of the parameters $\alpha_0, \alpha_1, \alpha_2, \dots, \alpha_k$ and as a result to two different model blends.

Implied volatility models are another important class of volatility models. The implied volatility is the value of the volatility parameter of a Black-Scholes option pricing equation that matches the theoretical prices of the options with the quoted market prices. A short description of

the Black-Scholes equation is given in Section 5.2. Let us assume that we have the market prices of call and put options for different maturities and all other parameters are known (except the volatility). We can estimate the volatility using the market prices by solving the reverse Black-Scholes problem. This volatility will be the implied volatility (IV). The IV is a function of the market price of the options, the underlying asset, the risk free rate, the exercise price, the time-to-expiration, and expected dividends. Unfortunately, there is no direct equation for computing the implied volatility from option prices (Hull, 2002), however, a search method can be introduced which allows us to compute the implied volatility with a good accuracy. One of the biggest challenges of this approach is a presence of special patterns (“smiles”) of the implied volatility. This issue will be addressed in detail in Section 3. The implied volatilities are heavily used in practical applications as an estimate of the conditional volatility forecast.

2.4 Applications of Extreme Value Theory to the market risk

Let us return to the discussion of the equations for the Value-at-Risk and the Expected-Shortfall (8.1) – (8.2). We have discussed different methods for conditional volatility estimation in the previous chapter. In this chapter we will discuss application of Extreme Value Theory (EVT) for modeling the distribution of residuals Z in equation (6). Let us assume that residuals Z have an unknown distribution function $F(x) = P\{X_i \leq x\}$. We are interested in the so-called excess distribution $F_u(y) = P\{X - u \leq y \mid X > u\}$, where $0 \leq y < x_0 - u, x_0 \leq \infty$. In case of market risk management, the excess distribution is interpreted as the probability of a loss that exceeds the threshold u . In general, the F is a distribution with an infinite right end, than theoretically we are exposed to the arbitrary large losses.

The unknown excess distribution F could be modeled by a number of different distributions, e.g., normal, t-distribution, etc. But there is another distribution that is more suitable for our purpose. A Generalized Pareto Distribution (GPD) is a continuous distribution of the following form:

$$G_{\xi, \beta}(x) = \begin{cases} 1 - \left(1 + \frac{\xi x}{\beta}\right)^{-\frac{1}{\xi}}, & \xi \neq 0, \\ 1 - e^{-\frac{x}{\beta}}, & \xi = 0, \end{cases} \quad (2.27)$$

where $\beta > 0$, and where $x \geq 0$ when $\xi \geq 0$ and $0 \leq x \leq -\beta/\xi$ when $\xi < 0$. The main result of the EVT is the limit theorem (McNeil A. , 1999). It says that, for a large class of underlying distributions F , the threshold u is progressively increasing; the excess distribution F_u converges to a Generalized Pareto Distribution G . The discussion of this theorem can be found in (McNeil A. , 1999). The ETV suggests a distribution that is a natural extension of wide variety of distributions (including the normal and the t-scale) on the extreme events.

We are interested in the procedure of estimating the parameters of the Generalized Pareto Distribution as well as explicit expressions for the VaR and the ES. Fortunately, EVT provides us this information. The tail estimate for the GPD is given by:

$$\hat{F}(x) = 1 - \frac{N_u}{n} \left(1 + \xi \frac{x - u}{\hat{\beta}}\right)^{-\frac{1}{\xi}}, \quad (2.28)$$

where

n is the total number of observations,

N_u is the number of observations exceeding threshold u .

The maximum log likelihood uses the tail estimate (2.28) to estimate the parameters of the GPD for given data. The EVT also gives the relationship for the Value-at-Risk for a given probability $q > F(u)$:

$$\widehat{VaR}_q = u + \frac{\hat{\beta}}{\hat{\xi}} \left(\left(\frac{n}{N_u} (1 - q) \right)^{-\hat{\xi}} - 1 \right). \quad (2.29)$$

The estimate of the Expected Shortfall can be obtained as follows:

$$\widehat{ES}_q = \frac{\widehat{VaR}_q}{1 - \hat{\xi}} + \frac{\hat{\beta} - \hat{\xi}u}{1 - \hat{\xi}}. \quad (2.30)$$

The EVT is applicable for problems of market risk management (McNeil A. , 1999). It suggests a special distribution for the tails of the distribution of losses and explicit equations for the key parameters. We will apply the results of the EVT for estimating the VaR and the ES at equation (2.9)-(2.10).

2.5 Comparison of the volatility forecasting models

The empirical tests of the volatility models were performed on various classes of assets. All of the models were tested on daily data. The description of the data sets and their characteristics are given in Table 2-1.

Table 2-1. The data sets with their characteristics.

N	Data set	Description	Number of instruments	Number of observations per instrument
1	Bonds	Returns of government bonds	5 bonds	2500
2	Commodities	Returns of commodity futures; including energy futures	44 futures	2500
3	FX	Returns of FOREX currency pairs	7 pairs	2500
4	Indices	Returns of major world indices	18 indices	2500
5	Stocks	Returns of individual stocks. The stocks are from main world indices.	267 stocks	1000 to 2500
6	Implied Volatility Indices	Returns of some stock indices and a daily implied volatility.	6 indices	less than 600
7	Implied Volatility Stocks	Returns of some stocks and their daily implied volatility.	6 stocks	less than 600

The 5-fold cross validation was used to divide each data set in training and testing subsets. The models were calibrated on the training subset and the resulting error was calculated over the testing subsets. The detailed description of a cross validation methods can be found in Section 5.3.

For each of the data sets the Root Mean Square Error (2.11) and the Mean Heteroscedastic Square Error (2.12) were computed. The Risk Characteristics were computed in order to fully compare the volatility forecasting methods. The following characteristics were computed:

1. The quantile $q_\alpha(Z)$ of the distribution of residuals Z in equation (2.6). This quantile is a constant which will give the Value-at-Risk (VaR) of the portfolio, after a multiplication by conditional volatility estimation. The quantile is estimated on the training set using (2.29).
2. The \widehat{ES}_q is the Expected Shortfall of the portfolio in terms of the VaR. It is computed by the following equation

$$\widehat{ES}_q = \frac{ES_q}{q_\alpha(Z)}, \quad (2.31)$$

where ES_q is computed with equation (2.30). This characteristic is computed on a training set.

3. The *Observed ES (OES)* is an empirically computed average value of the volatility that exceeds the VaR threshold. This characteristic is computed on the testing set. The \widehat{ES}_q is expected to be equal to the *OES*.
4. The *Confidence level (CL)* is a percentage of the observations, for which the absolute value of returns exceed the VaR threshold. The *CL* is computed on the testing set and is expected to be equal to $\alpha = 99.8\%$.

We have calculated an average value of the error functions and Risk Characteristics on all of the datasets. We can divide all of the datasets into two types based on the presence of the implied volatility data. The first type includes the datasets without the implied volatility: Bonds, Commodities, FX, Indices and Stocks data sets. The average results over these datasets are given in Table 2-2.

Table 2-2. Performance of the volatility forecasting models on data sets, that do not include the implied volatility data.

N	Characteristic	Method					
		EWMA (λ^*)	GARCH (1,1)	ARCH(1)	EGARCH (1,1)	MB (RMSE)	MB (MHSE)
1	RMSE	1.3790	1.5083	1.6361	1.4980	1.2555	1.8751
2	MHSE	0.6413	1.6618	0.6422	5.2793	1.1081	0.4682
3	$q_\alpha(Z)$	4.7075	8.5885	4.5840	11.453	6.2752	2.9838
4	\widehat{ES}_q	1.3987	1.4216	1.3940	1.8593	1.3821	1.3605
5	OES	1.3429	1.4206	1.3824	1.7887	1.3386	1.3620
6	CL	99.8035	99.7359	99.7275	99.6902	99.7942	99.8069

The second type of data sets includes values of the implied volatility data. The volatility of the 45-days at-the-money options is taken as a value of the implied volatility. The results of the volatility forecasting methods applied on a these datasets are given in Table 2-3. However, we should note that the results found in Table 2-3 are less trustworthy, than those of the Table 2-2. This is because there are much fewer observations in the datasets with the implied volatility, than without.

Table 2-3. Comparison of performance of the volatility forecasting models with the implied volatility.

N	Characteristic	Method						
		EWMA (λ^*)	GARCH (1,1)	ARCH (1)	EGARCH (1,1)	MB (RMSE)	MB (MHSE)	IV
1	RMSE	2.3634	2.4070	2.7118	2.2735	2.1147	3.4855	2.1319
2	MHSE	0.9216	1.3698	0.7413	2.1329	1.4856	0.6135	0.8674
3	$q_\alpha(Z)$	4.3867	6.2613	4.4517	8.1206	6.2821	2.5313	4.2425
4	\bar{ES}_q	1.9195	1.5197	1.4408	1.7887	1.9493	1.5512	1.6859
5	OES	1.0045	1.0341	1.0317	1.3096	0.8283	1.1624	1.1235
6	CL	99.8391	99.7501	99.7555	99.7224	99.8556	99.7998	99.7696

The optimal parameter of EWMA λ^* was determined using the maximum likelihood function of a t-Distribution. The values of λ^* were computed separately for each of the datasets.

The EWMA is a simple but effective method for conditional volatility estimation. It gives a good estimate of the volatility in terms of error functions, as well as an accurate estimate of the Risk Characteristics. The deviation between the theoretical Expected Shortfall and the observed Expected Shortfall is relatively small. The EWMA method is easy to implement and does not require frequent recalibration of its parameters.

The Heteroscedastic family of models, however, did not give superior results to simple models. All models of this family (except ARCH) give poor results in terms of both the RMSE and the MHSE error functions. The ARCH/GARCH and EGARCH models give an accurate estimate of the Expected shortfall. This family of methods has a significantly lower performance than EWMA over all characteristics. Although, heteroscedastic models are interesting from a theoretical point of view, their use for purposes of risk management is inappropriate.

We have considered two different Model Blending techniques. The first one is designed to minimize the RMSE error function, while the second is designed to minimize the MHSE error. The MB methods give the best accuracy in terms of error functions: the MB (RMSE) gives the lowest RMSE and the MB (MHSE) gives the lowest MHSE. Both of these methods give a good estimate of the Expected Shortfall. The MB (MHSE) gives the lowest Mean Benchmark Deviation among other volatility forecasting methods. It means that the MB (MHSE) gives the best conditional volatility estimate, keeping the values of the Risk Characteristics the same as the method used in the Correlation Haircut system. A good performance of the Model Blending approach based on the MHSE

error function has shown that, the MHSE error has an advantage over the RMSE for the problems of the conditional volatility estimation. We can conclude that a method which will give a low MHSE error most probably will also produce suitable values of the Risk Characteristics. The main disadvantage of the Model Blending approach is that it requires the implementation and calibration not of one model but of all the models used in blending. The parameter calibration procedure is complicated and could potentially require repeated recalibration of the parameters. The MB method is a “black-box” approach; the forecasts made by this method are often non-intuitive.

We have compared the results of the Historical volatility models with the Implied Volatility approach. The results are represented in Table 2-3. We would like to stress that those results are less reliable, because the comparison was made only for a limited number of financial products. For the selected products the IV gives low values of error functions as well as a good estimate of the Risk Characteristics. The IV estimates volatility based on the market values of the options. This connection with the market makes IV an attractive technique. The main disadvantage of this method is that on average it tends to overestimate the observed volatility. Implied volatility reflects the ‘fears’ of market players. Another disadvantage is that for some assets this method is hard to implement as there are no options traded or the traded options are not liquid.

We have compared several popular methods for the conditional volatility estimation. Each method has its advantages and disadvantages, which were described in this chapter. Some methods are simple but yield poor results, while other methods provide improved results but are difficult to implement. In short there is no perfect approach. The Exponentially Weighted Moving Average method with the optimal set of parameters provides good results in terms of both error functions and Risk Characteristics. Although some methods (MB-MHSE) outperform EWMA, the EWMA method is easy to implement and to calibrate. The Implied Volatility can be also successfully used as a volatility estimation technique for the liquid assets. We can conclude that the combination of EWMA and IV approach should be used for purposes of risk management.

3 Modeling implied volatility surfaces

3.1 Risk of volatility “smile” and its impact

European options are often priced or hedged using the Black-Scholes model (Black & Scholes, 1973). The model gives an one-to-one relationship between the price of the European put or call option and the implied volatility. It assumes that the price of the underlying asset follows the geometric Brownian motion with constant volatility. This is a rather crude assumption, because it implies that the volatility parameter is equal for all the strikes. The Black-Scholes conditions never hold exactly in the markets. This happens due to different factors, for example, jumps in underlying asset prices, movements of volatility over time, transaction costs, etc. Therefore, practitioners often use different values of implied volatility for different strike prices. This forms a specific pattern, called the “implied volatility smile”. Although, this pattern could be in a form of a “skew”, “smile” or a “sneer”, we will refer to it as a “skew”. The implied volatility surface is a more general representation of implied volatility smile pattern. By the implied volatility surface we will understand the dependence between the price of the underlying asset, strike price of the option, time to maturity of the option and its implied volatility.

Changes of the implied volatility have a significant influence on the value of the option position. An incorrect estimate of the implied volatility and its expected shifts could lead to a significant miss-pricing of the options. That is why modeling the implied volatility surfaces plays an important role in financial risk management. In this section we will test several models for implied volatility surface modeling. We will include different types of polynomial fitting as well as a stochastic volatility models in our analysis.

3.2 Representation of the moneyness

The volatility skew or surface is usually expressed in terms of moneyness X rather than in terms of a simple strike price K . Moneyness is a ratio between the strike price K and the price of the underlying asset S . We will focus on two different representations of moneyness. The first representation of moneyness is given by the following equation:

$$X = \ln(K/S). \quad (3.1)$$

Alternatively, we can formulate moneyness in terms of time to expiration. Let us denote time to expiration as $\tau = T - t$, where T is a date of expiration of a given option and t is a current date. Then the moneyness can be redefined as:

$$X = \frac{\ln(K/S)}{\sqrt{\tau}}. \quad (3.2)$$

An option is said to be at-the-money if $X = 0$. A call (put) option is said to be in-the-money (out-of-the-money) for $X < 0$ and out-of-the-money (in-the-money) for $X > 0$. The comparison of the volatility skew for different moneyness is represented in Figure 3-1.




Figure 3-1: The observed implied volatility of the AEX index options on 24-10-2007. Left: representation in the log-moneyness (3.1). Right: representation in the time adjusted moneyness (3.2). Different colors represent different time-to-expiration in days.

The implied volatility surface is the representation of the implied volatility as a function of moneyness and time to expiration $\sigma_B = \sigma_B(X, \tau)$. An example of the volatility surface is given in Figure 3-2.




Figure 3-2: An example of the implied volatility surface of AEX index options on 26-01-2009.

3.3 Models of the implied volatility surface

There are a number of mathematical models that describe the implied volatility surfaces. An overview of some of them will be given in this chapter. We will focus on the models which are extensively described in the literature and have a proven record of effectiveness.

We will start our overview with the polynomial models. The **Cubic model** is a model for volatility surfaces. It uses the time adjusted form of moneyness (3.2). This model treats the implied volatility as a cubic function of moneyness X and a quadratic function of the time to expiration τ . The model is described by the following equation:

$$\sigma_{impl} = a_0 + a_1X + a_2X^2 + a_3X^3 + a_4\tau + a_5\tau^2, \quad (3.3)$$

Parameters a_0, a_1, a_2, a_3, a_4 and a_5 are estimated using the least squares method. The Cubic model describes whole volatility surface with one equation.




Figure 3-3: An example of the fit of the Cubic model to the AEX index options implied volatility surface on 26-01-

2009.

So for any given moneyness X and time to expiration τ one can find the value of implied volatility. This model is a smooth function of moneyness and time to expiry. An example of the fit of the cubic surface model is given in **Error! Reference source not found.**

With the analogies to the Cubic model, we can build a **Spline model** of the volatility surface. We are using the time adjusted form of moneyness (3.2). Let us introduce the dummy variable D :

$$D = \begin{cases} 0, & X < 0, \\ 1, & X \geq 0. \end{cases} \quad (3.4)$$

The volatility function is given by:

$$\begin{aligned} \sigma_{impl} = & a_0 + a_1X + a_2X^2 + a_3\tau + a_4\tau^2 \\ & + D(a_5 + a_6X + a_7X^2 + a_8\tau + a_9\tau^2). \end{aligned} \quad (3.5)$$

We would like the function of volatility to be continuous and differentiable. This is achieved by adding the following constrains:

$$a_5 + a_6 \cdot 0 + a_7 \cdot 0^2 + a_8\tau + a_9\tau^2 = 0, \quad (3.6)$$

$$Da_6 = 0. \quad (3.7)$$

We get the following function of the implied volatility, after rearranging the terms of (3.5):

$$\sigma_{impl} = a_0 + a_1X + a_2X^2 + a_3\tau + a_4\tau^2 + Da_7X^2.. \quad (3.8)$$

The parameters a_0, a_1, a_2, a_3, a_4 and a_7 are fitted using the linear least squares regression analysis. The whole volatility surface is described by six parameters. The Spline model gives a value of the volatility for any value of τ , and does not require any additional interpolation over the time to expiration. An example of the fit of this model is given in Figure 3-4.




Figure 3-4: An example of the fit of the Spline model to the volatility surface of AEX index options on 26-11-2008.

The **Stochastic alpha, beta, rho (SABR)** is another important class of volatility models. The SABR model was introduced in 2002 (Hagan, Kumar, Lesniewski, & Woodward, 2002). This model assumes some behavior of the underlying asset and connections with the values of the implied volatility. The shape of the volatility skew is derived analytically from these assumptions. Let us denote the today's forward price of the underlying asset by f and the forward price of the asset for a forward contract that matures on the fixed settlement date by $\hat{F}(t)$. Today's forward price is defined as $f = \hat{F}(0)$. The strike price of an European option is denoted by K . The forward price and the volatility are described by the following processes:

$$d\hat{F} = \hat{\alpha}\hat{F}^\beta dW_1, \quad \hat{F}(0) = f, \tag{3.9}$$

$$d\hat{\alpha} = v\hat{\alpha}dW_2, \quad \hat{\alpha}(0) = \alpha, \tag{3.8}$$

under the forward measure, where the two processes are correlated by :

$$dW_1 dW_2 = \rho, \quad (3.11)$$

where W_1 and W_2 are correlated Brownian motions. The equation for the implied volatility $\sigma_B(f, K)$ is derived from (3.9)-(3.11) to satisfy Black's equation:

$$V_{call} = D(T)\{f\mathcal{N}(d_1) - KN(d_2)\}, \quad (3.92)$$

$$V_{put} = V_{call} + D(T)[K - f], \quad (3.103)$$

$$d_{1,2} = \frac{\log(f/K) \pm \frac{1}{2}\sigma_B^2\tau}{\sigma_B\sqrt{\tau}}, \quad (3.14)$$

where,

V_{call}, V_{put} are values of European call and put option,

$D(t)$ is a discount factor at day t ,

$\mathcal{N}(\cdot)$ is the cumulative probability distribution function for the standard normal distribution.

The implied volatility is given by

$$\sigma_B(K, f) = \frac{\alpha}{(fK)^{\frac{1-\beta}{2}} \left\{ 1 + \frac{(1-\beta)^2}{24} \log^2 f/K + \frac{(1-\beta)^4}{1920} \log^4 f/K + \dots \right\}} \left(\frac{z}{x(z)} \right) \cdot \left\{ 1 + \left[\frac{(1-\beta)^2}{24} \frac{\alpha^2}{f^{2-2\beta}} + \frac{1}{4} \frac{\rho\beta\nu\alpha}{(fK)^{(1-\beta)/2}} + \frac{2-3\rho^2}{24} \nu^2 \right] \tau + \dots \right\} \quad (3.15)$$

Here

$$z = \frac{\nu}{\alpha} (fK)^{\frac{1-\beta}{2}} \log f/K \quad (3.16)$$

and $x(z)$ is defined by

$$x(z) = \log \left\{ \frac{\sqrt{1 - 2\rho z + z^2} + z - \rho}{1 - \rho} \right\}. \quad (3.17)$$

For the special case of at-the-money options, options struck at $K = f$ (or $X = 0$), this equation reduces to :

$$\sigma_B(f, f) = \frac{\alpha}{(fK)^{\frac{1-\beta}{2}}} \cdot \left\{ 1 + \left[\frac{(1-\beta)^2}{24} \frac{\alpha^2}{f^{2-2\beta}} + \frac{1}{4} \frac{\rho\beta\nu\alpha}{(fK)^{(1-\beta)/2}} + \frac{2-3\rho^2}{24} \nu^2 \right] \tau + \dots \right\} \quad (3.18)$$

The detailed description and derivation of equations (3.15)-(3.18) can be found in (Hagan, Kumar, Lesniewski, & Woodward, 2002). The SABR model can be easily rewritten to take relative strikes (moneyness) as input. In this case:

$$\sigma_B\left(\frac{K}{f}, 1\right) = \sigma_B(K, f). \quad (3.19)$$

There is a special procedure for calibrating the parameters of the SABR model. We will start with the β parameter. (Hagan, Kumar, Lesniewski, & Woodward, 2002) and (Poon S. , 2008) suggest to fix the β parameter in advance and not to change it. There are two special cases: $\beta = 0$ and $\beta = 1$. The first one represents a stochastic normal model, while the second one ($\beta = 1$) represents a stochastic log-normal model. The value of β should be chosen to be between 0 and 1. We will focus on the case of $\beta = 1$. Once β is fixed, all other parameters can be adjusted to the value of the β . The particular value of β has little impact on the shape of the implied volatility skew. This parameter can be successfully chosen from “esthetical” considerations (West, 2005).

The α parameter refers to the value of the at-the-money volatility. There are two ways of estimating this parameter. It could be taken as a value of at-the-money volatility $\alpha = \sigma_{ATM}$. Or it could be estimated as a smallest positive solution of the following equation:

$$\frac{(1-\beta)^2}{24} \frac{\tau\alpha^3}{f^{2-2\beta}} + \frac{1}{4} \frac{\rho\beta\nu\tau\alpha^2}{f^{1-\beta}} + \left(1 + \frac{2-3\rho^2}{24} \nu^2\tau\right)\alpha - \sigma_{ATM}f^{1-\beta} = 0. \quad (3.11)$$

The α parameter should be chosen to be positive, $\alpha > 0$. Changes of the α parameter shift the implied volatility skew across the volatility axis. It controls the level of the implied volatility curve or surface. The parameters ν and ρ are calibrated to minimize the error between the observed implied volatilities and the SABR model (3.15)-(3.17). The ν parameter refers to the volatility of the implied volatility and is called ‘volvol’. The ρ parameter is a correlation coefficient between the at-the-money volatility and movements of the underlying asset. To calibrate ν and ρ the two dimensional Nelder-Mead algorithm was suggested by (West, 2005). The algorithm gives the optimal values ν^* and ρ^* that minimize the quadratic error:

$$(\nu^*, \rho^*) = \arg \min_{\substack{\nu > 0 \\ -1 \leq \rho \leq 1}} \sum_{i=1}^N (\sigma_{M,i} - \sigma_{B,i})^2, \quad (3.12)$$

where,

$\sigma_{M,i} \ i = 1, \dots, N$ are the values of the implied volatility obtained from the market prices of the options,

$\sigma_{B,i} \ i = 1, \dots, N$ are the SABR estimations of the implied volatility.

Now, we can summarize the procedure of fitting the parameters of the SABR model. Firstly, fix the value of β parameter. We are taking $\beta = 1$ in all of our experiments. Secondly, find the value of α . This can be done by solving (3.11) or by fixing $\alpha = \sigma_{ATM}$. For practical reasons, we will fix α to be equal to the volatility of at-the-money options. At last, the parameters ν and ρ are calibrated using procedure (3.12). All the remaining parameters of the SABR equation (3.15)-(3.17) are known for a given contract. Although, (3.15)-(3.17) looks complicated, it involves only simple mathematical operations and can be implemented rather easily.

The SABR model can be used as a model for a whole volatility surface or for the skew (fixed τ). We will discuss these two approaches separately. Under the first approach, the parameters $\alpha, \beta, \rho,$ and ν are calibrated for all given times to expiration $\tau_i, i = 1, \dots, l$. A point on the volatility surface is obtained by applying (3.15)-(3.17). An example of the SABR volatility is given in Figure 3-.




Figure 3-5: An example of the fit of the SABR model to the volatility surface of AEX index options on 9-7-2008.

A second approach is to fit a SABR skew for each observed time to expiration. And then interpolate the values of implied volatility for any arbitrary τ . The SABR parameters $\alpha_i, \beta_i, \rho_i$, and ν_i are calculated separately for each time to expiration $\tau_i, i = 1, \dots, l$. The implied volatility surface is built as a linear approximation of separate skews.

$$\sigma_B'(K, f, \tau') = \sigma_{B1}(K, f, \tau_1) \left(\frac{\tau_2 - \tau'}{\tau_2 - \tau_1} \right) + \sigma_{B2}(K, f, \tau_2) \left(\frac{\tau' - \tau_1}{\tau_2 - \tau_1} \right). \quad (3.13)$$

where,

σ_B' is the value of the implied volatility for any arbitrary time to expiration τ'
 σ_{B1}, σ_{B2} are values of the implied volatility calculated from time to expiration that are observed on the market τ_1, τ_2 .

We will refer to this method as **piecewise SABR (PSABR)**. An example of the fit of the PSABR is given in Figure 3-




Figure 3-6: An example of the fit of the PSABR model to the observed implied volatility skew of the AEX index options on 17-3-2008, with time-to-expiration of 95 days.

3.4 Modeling the dynamics of volatility surfaces

The value of the implied volatilities changes with time, deforming the shape of the implied volatility surface. The evolution in time of this surface captures the changes in the options market. While a model with a large number of parameters may calibrate well the volatility surface on a given day, the same model parameters may give poor result on the next day. Any risk management system tries to estimate the future (short term forecast) behavior of the volatility surface. Modeling the dynamics of the implied volatility surface is an important task from practical point of view. In this chapter we will discuss different techniques to model the dynamics of the implied volatility surfaces. We will focus on two different approaches. One will be applicable to the cubic and the spline model; the other will be used for SABR models.

The Cubic (3.3) and the Spline (3.8) models of the volatility surface use parameters $a_i, i = 1, \dots, n$ to model the shape of the surface. The dynamics of the surface is treated as dynamics of these parameters. So, we can assume that $a_i = a_i(t), i = 1, \dots, n$. In order to reduce the dimensionality of the problem, we apply Principal Component Analysis (PCA) to the values of $a_i, i = 1, \dots, n$. The PCA is a statistical technique widely used in practice as a preprocessing technique. The details of the PCA could be found in (Shlens, 2005). Let us denote by matrix A ; the $T \times n$ matrix of observations of coefficients $a_i(t), i = 1, \dots, n; t = 1, \dots, T$ of the implied volatility surface. After applying PCA to the matrix A , we will obtain the matrixes P, C and a vector l . P is a $n \times n$ matrix, each column containing coefficients for one principal component. The columns are in order of decreasing component variance. C is an $T \times n$ matrix, the representation of A in the principal component space. l is a vector containing the eigenvalues of the covariance matrix of A . An example of l is given on Figure 3-. PCA is theoretically the optimal linear scheme, in terms of least mean squares error, for compressing a set of highly-dimensional vectors to a set of lower-dimensional vectors and then reconstructing the original set (Shlens, 2005). Working with coefficients of the models (3.3), (3.8) can be substituted by working with the matrix C . Moreover, the first few principal components explain most of the variance of the A . For practical considerations we will focus on the first two principal components. The dynamics of the volatility surface over time is explained by dynamics of these two variables. An example of the dynamics of the first and the second principal component is given in Figure 3-.




Figure 3-7: A PCA of the coefficients of the Cubic model of the fit of AEX index implied volatility surface. The first two components explain almost all variance of the coefficients.




Figure 3-8: The dynamics of the 1st and the 2nd principal component of the coefficients of the Cubic model for the AEX index implied volatility.

We assume that the two first principal components $c_1(t)$ and $c_2(t)$ have a significant value of sample partial autocorrelation. A sample partial autocorrelation function for the first two principal components is given in Figure 3-.




Figure 3-9: A sample partial autocorrelation function of the 1st and the 2nd principal components of the Cubic model applied to the AEX index implied volatility surface. Both, the 1st and the 2nd principal component exhibit a significant value of the autocorrelation for a 1 day lag.

The dynamics of $c_1(t)$ and $c_2(t)$ can be modeled with an Autoregressive moving average model (ARMA):

$$\hat{c}(t+1) = \sum_{i=1}^p \mu_i c(t+1-i) + \sum_{j=1}^q \gamma_j \xi_{t+1-j}, \quad (3.14)$$

where $\mu_1, \mu_2, \dots, \mu_p$ and $\gamma_1, \gamma_2, \dots, \gamma_q$ are parameters of the model and $\xi_1, \xi_2, \dots, \xi_q$ are the error terms of the model $c(t) = \hat{c}(t) + \xi_t$. We will refer to this model as the $ARMA(p, q)$ model. For a particular case of modeling $c_1(t)$ and $c_2(t)$, we will take $p = 1$ and $q = 1$. The parameters μ_1 and γ_1 are estimated using a least squares method.

Now, we can summarize the procedure of modeling the dynamics of the implied volatility surface for cubic and spline models. First, we apply the PCA to the historical observations of $a_i(t), i = 1, \dots, n$, to obtain observations of $c_1(t)$ and $c_2(t)$. Then the ARMA model (3.14) is calibrated and applied to get the forecast $\hat{c}_1(t+1)$ and $\hat{c}_2(t+1)$. After that, the PCA is applied “backwards”

(Shlens, 2005) to construct the forecast of $\hat{\alpha}_i(t+1), i = 1, \dots, n$. The forecasted surface is constructed from $\hat{\alpha}_i(t+1), i = 1, \dots, n$ using **Error! Reference source not found.** or (3.8). The Principal Component Analysis is a popular statistical technique to reduce the dimensionality of the problem. We are using PCA in a somewhat nonstandard way. We are reducing the relatively small number of variables (coefficients of a Spline or Cubic model) even more. The main reason for the PCA for our problem is to switch to another space of the no-correlated factors, that fully describe the dynamic of the implied volatility surface.

Apart from the previously discussed model of dynamics, the SABR model already assumes certain dynamics of the volatility and the underlying asset. This dynamics is expressed by a system of stochastic differential equations:

$$d\hat{F} = \hat{\alpha}\hat{F}^\beta dW_1, \quad \hat{F}(0) = f, \quad (3.24)$$

$$d\hat{\alpha} = v\hat{\alpha}dW_2, \quad \hat{\alpha}(0) = \alpha, \quad (3.25)$$

where W_1 and W_2 are correlated Brownian motions

$$dW_1dW_2 = \rho. \quad (3.26)$$

Instead of the forward price of the underlying asset f , the current spot price of the underlying asset S can be used. Suppose, the SABR model is already calibrated and parameters α, β, ρ, v are already known. Then (3.24)-(3.26) can be simulated with the Monte-Carlo method (MC). (3.24)-(3.26) are model dependent on random variables (changes of Brownian motion). Under the MC method we assume some distribution of these random variables and generate N_{MC} realizations of them. Based on these realizations we calculate N_{MC} outputs of the model. The average of the outputs is used as a forecast of the process. In our particular case, the change of Brownian motion has a normal distribution $dW_i \sim N(0,1), i = 1,2$. We will use the following approach in order to generate correlated random numbers dW_1 and dW_2 with a correlation coefficient ρ . First, we generate two uncorrelated sequences of normal random numbers dW_1, Z . The correlated sequence of random numbers is constructed in the following way:

$$dW_2 = \rho dW_1 + \sqrt{1 - \rho^2}Z. \quad (3.27)$$

The sample paths of the underlying asset \hat{F} and volatility-like parameter $\hat{\alpha}$ are generated from (3.24) and (3.25). The SABR equations (3.15)-(3.17) are applied for each path of the MC simulation. This

results to the N_{MC} simulated implied volatility surfaces. The forecast of the implied volatility surface is an average surface over the simulated paths.

We should note that the forecasting procedure is slightly different for the SABR surface model than for the PSABR model. The SABR surface is described by 4 parameters α, β, ρ and ν . The forecasting algorithm is the same as described earlier. But the PSABR model is described by $4 \times N_\tau$ parameters, where N_τ is a number of is is different time to expiry τ in the observed option portfolio. The dynamics of each volatility skew (fixed τ) is considered to be a separate process. The MC simulation is applied separately for each of the N_τ sets of parameters. This will lead to the N_τ forecasted implied volatility skews. The implied volatility surface is constructed from these skews by applying (3.13).

The Sticky-Strike rule is another approach which is used for estimating the shifts of the implied volatility. The main idea is the assumption that if the price of the underlying asset changes, than the implied volatility of an option with a given strike does not change. The Sticky-Strike rule is very popular among practitioners, because it easy to understand and does not require complex computations. Let us formalize this approach. The Sticky-Strike rule suggests that:

$$\sigma(K, f(t), \tau(t)) = \sigma(K, f(t-1), \tau(t-1)) + \epsilon, \quad (3.28)$$

where

$\sigma(K, f(t-1), \tau(t-1))$ is the value of the implied volatility on a day $t-1$,

$\sigma(K, f(t), \tau(t))$ is the value of the implied volatility on a day t , for the same strike K .

The Sticky-Strike rule is not a model of the implied volatility surface. It does not provide an equation which can give a value of volatility for any strike, moneyness or time to expiration. It is only an empirical rule to estimate the future behavior of the implied volatility of the options in a given portfolio.

3.5 Comparison of the volatility surface models

The models and methods described in the previous chapters were tested on market data, in order to determine their advantages and disadvantages. The testing data consists of 10 separate sets, each corresponding to a different underlying asset. The list of data sets with a short description is given in Table 3-1.

Table 3-1. The characteristics of the data sets used for the analysis.

N	Underlying symbol	Description	Dates of observations	Number of days	Adjusted for implied dividends (Y/N)
1	AEX	Amsterdam Stock Index	7-6-2007 to 1-4-2009	467	Y
2	DAX	German Stock Index	7-6-2007 to 11-5-2009	496	Y
3	FTSE	UK Stock Index	7-6-2007 to 11-5-2009	494	Y
4	EURSTOX	European Stock Index	7-6-2007 to 11-5-2009	494	Y
5	S&P 500	US Stock Index	7-6-2007 to 11-5-2009	486	Y
6	XJO	Australian Stock Index	7-6-2007 to 11-5-2009	499	N
7	DBK	Deutsche Bank Stock	7-6-2007 to 11-5-2009	494	N
8	VOW	Volkswagen Stock	7-6-2007 to 11-5-2009	494	N
9	ALV	Allianz Stock	7-6-2007 to 11-5-2009	494	N
10	ADS	Adidas Stock	7-6-2007 to 11-5-2009	494	N

The data set includes the values of implied volatilities of put and call options with different time to expiration. Each record is characterized by date of trade, strike price, price of the underlying asset, time to expiration, implied volatility (derived from the market price of the option) and type of the option (put or call). Put –call parity (Hull, 2002) suggests that implied volatilities of call options should not significantly deviate from the implied volatilities of the put options for the same moneyness. However, in practical applications this deviation is commonly observed. The adjustment for implied dividends of the underlying asset can be made, in order to eliminate these deviations (Hafner & Wallmeier, 2000). Part of the data set is adjusted for the value of the implied dividends. As a result, the values of the implied volatility for call and put options are almost equal. This adjustment is mainly done when the underlying asset is a stock index (see Table 3-1). The other part of the data is not adjusted for the value of implied dividends. An example of the implied volatility skew with and without adjustment is given in Figure 3-3.




Figure 3-3: An example of the observed implied volatility for the datasets adjusted and not adjusted for the value of implied dividends. Left: implied volatility of the AEX index options on 1-23-2009. Right: implied volatility surface of options on the Volkswagen (VOW) shares on 1-23-2009.

Data cleaning should be applied before testing the methods. Data cleaning is a process of removing inaccurate or unreliable records from the data set. We will remove data with a very small or a big value of delta. The options delta is a sensitivity of a value of the option to the changes of the underlying asset:

$$\Delta = \frac{\partial V}{\partial S}. \quad (3.29)$$

We will remove records from the data set with $|\Delta| > 0.9$ and $|\Delta| < 0.01$. Options that are close to maturity exhibit significant jumps in volatility. These options are very sensitive to the changes of moneyness. We will exclude from the data set observations of options with time to expiration less than 60 days. Far out-of-the-money options can have potentially an inaccurate estimate of the implied volatility. Options with moneyness around zero have the most accurate estimate of implied volatility. We will remove the options with an absolute value of moneyness greater than five, $|X| > 5$. By this we will remove only significant outliers.

In order to estimate the forecasting performance and quality of model fit of some methods or to compare several methods we should define the error function. The observations of the implied volatility are not equally trusted. Some options have higher sensitivity of the volatility to the changes of underlying asset, while others have a lower sensitivity. We will build an error measure that reflects this issue. The error measure is a weighted modification of the square error. These weights will

reflect the sensitivity. Let us denote by ν the vega and by Δ the delta of the option. The weight will be given by the vega to delta ratio:

$$w = \left| \frac{\nu}{\Delta} \right|. \quad (3.15)$$

Low values of w indicate a high implied volatility sensitivity to the changes in the underlying price. This should be reflected in the small weight of the error. High values indicate low implied volatility sensitivities with respect to underlying price movements. Therefore the weight of the error should be large. The error function is given by the Weighted Mean Square Error:

$$WMSE = \frac{1}{N} \sum_{i=1}^N w_i \left(\hat{\sigma}_{B,i}(X, \tau) - \sigma_{B,i}(X, \tau) \right)^2, \quad (3.16)$$

where

$\sigma_B(X, \tau)$ is the value of the implied volatility observed on the market,

$\hat{\sigma}_B(X, \tau)$ is the value of the implied volatility estimated by one of the volatility surface models for the same time to expiration and moneyness,

w_i are the weights of the i – th observation, calculated with the sensitivity ratio,

N is the number of observations in a data set over all trading days

The error function (3.16) calculates an average quadratic deviation between the observed and the modeled data, and gives more weight to the “more trusted” observations. This error function is used to calibrate the parameters of all the models, as well as to calculate the error for model comparison. We should note, that for technical reasons the sensitivity ratio (3.15) is normalized from $[0; 1]$ on a daily basis.

The WMSE is calculated for 5 different tests. The quality of fit is the first one. It is characterized by the RMSE between the implied volatility estimated by models and observed volatility on the same trading day. The forecasting power of the models is tested by the next four tests. We will apply dynamic models to build 1 and 5 day forecasts of the volatility skews. We will refer to these tests as tests of dynamics. The “rolling horizon” technique is used for these tests. Let us suppose that we want to build a forecast of the skew for day t , then we will use observations from days $t - N_{tr}$ to $t - 1$ to calibrate the dynamic models. Then for day $t + 1$ the “horizon” is “moved”, so that days $t - N_{tr} + 1$ to t are used for calibration. An equivalent technique is used for the five

days ahead forecast. In practical applications we will take $N_{tr} = 100$. The remaining two tests are applied to check how the models can “hold” the volatility skew pattern. The WMSE is calculated between the model results and observed implied volatility on day t , similar to the quality of fit test. But, in this case the parameters of the models are calibrated on observations of day $t - 1$ or $t - 5$. We will refer to these tests as “static tests”. The averaged WMSE of the models over data sets adjusted for the value of implied dividends is given in Table 3-2. The results of the test for the remaining data sets is given in Table 3-3. The WMSE for each of the underlying assets is given in section 5.5.

Table 3-2. Averaged results of the tests for the following underlying assets: AEX, DAX, FTSE, EUROSTOXX, and S&P500.

N	Model	Quality of fit	Static models		Dynamic models	
			1 day forecast	5 days forecast	1 day forecast	5 days forecast
1	Cubic	0.82596	2.44567	5.37869	2.78673	7.89823
2	Spline	0.63001	1.97957	4.65466	2.53439	6.47271
3	SABR	4.22855	5.11622	7.82890	5.11794	7.32580
4	PSABR	2.32145	3.27966	5.88138	4.50680	6.70848
5	Sticky Strike	N/A	1.23520	3.02228	N/A	N/A

Table 3-3. Averaged results of the tests for the following underlying assets: XJO, DBK, VOW, ALV and ADS.

N	Model	Quality of fit	Static models		Dynamic models	
			1 day forecast	5 days forecast	1 day forecast	5 days forecast
1	Cubic	4.16422	10.60288	22.37566	18.72404	50.34940
2	Spline	3.94045	9.54097	20.81255	12.61659	34.76029
3	SABR	14.20002	21.10567	38.06475	21.11537	33.90157
4	PSABR	5.47680	11.14137	23.03782	13.04728	23.03009
5	Sticky Strike	N/A	7.42659	16.40627	N/A	N/A

We can shortly summarize the advantages of each model, given the results of the tests. The Cubic and Spline models approximate the implied volatility surface with the function of a certain form. On average, the Spline model performs better than the Cubic model. Good performance of the dynamic version of the Spline model is an empirical evidence of the dependence of the dynamics of the surface of two principal components. Both of these models use much less parameters, than the PSABR model. Both of these models use much less parameters, than the PSABR model. Spline and Cubic model use one set of parameters to approximate whole surface, while PSABR use a separate set of parameters for each skew (which corresponding to fixed time to expiry).

The SABR and the PSABR assume a certain model for the joint dynamics of the volatility and the underlying asset. Unfortunately, the SABR model gives a higher error than other models. The PSABR model gives much better results. Although, the PSABR does not give the best forecast, it models the skew rather effectively, in case of insufficient or “bad” data. The SABR family of models assumes some shape (“skew”, “smile”, “sneer”, etc.) of the volatility skew and fits it to the given data. The SABR gives a more “theoretical” shape of the skew. This results to a higher fitting error, but lower relative forecasting error (error of forecast divided by the error of fit). Another advantage of the SABR model is that each parameter has a certain meaning. The ρ parameter is a correlation between the changes of at-the-money volatility and the underlying asset, ν is so-called “volvol” the volatility of the volatility, and α is associated with at-the-money volatility. These values are a useful “complementary” product of the SABR model, and can be used in other applications of risk management.

The Sticky Strike rule is a simple, but very effective way to estimate the shifts of the implied volatility. The WMSE of this method is one of the lowest over all data sets. The main disadvantage of this method is that it estimates the implied volatility only of those options that are being continuously tracked. In practical applications, however, not all the options are tracked to compute the implied volatility. Usually, the limited portfolio is considered; it makes it impossible to apply the Sticky Strike rule for any strike, moneyness and time to expiry. The sticky Strike is simple method; it does not require any computations or parameter estimates and can be effectively used as a complementary method.

Modeling the implied volatility surface is a challenging task. In this work we have tried several methods and approaches to solve this problem. We have developed and tested a number of models. We have demonstrated that there is no single model, which significantly outperforms the other. Each model has its advantages and disadvantages. The Spline model is perhaps, the most effective to minimize the fitting error. The SABR model has a set of meaningful parameters and a more significant “forecasting power” It performs better on an incomplete or missing data.

4 Conclusions and practical recommendations

This project is dedicated to addressing the problem of volatility modeling in financial markets. We focused on two major aspects of the volatility problem: the conditional volatility estimation and modeling volatility surfaces. The volatility modeling problem was studied with the purposes of financial risk management.

The first part of this thesis deals with the problem of conditional volatility estimation. We have selected several methods that are heavily used in practice and tested their accuracy using a number of different classes of real data. Each family of methods has its advantages and disadvantages, which are described in this work. Some methods yield poor results (e.g., the heteroscedastic family of models), while the others provide improved results but are difficult to implement (e.g., models blending). In short, there is no single perfect approach. Nevertheless, we found that the Exponentially Weighted Moving Average method is efficient and is relatively easy to implement. We have described the procedure to calibrate the parameters of this model. We have also tested Model Blending techniques. This is a relatively new approach, and we confirmed that it can be successfully used for volatility forecasting. The Model Blending approach certainly provides a superior accuracy over other methods. Its practical application, however, is compromised by an extremely complex procedure of parameter calibration. Given all these considerations, we suggest the following recommendations for conditional volatility estimation: a combination of the EWMA model (with properly calibrated parameters) and the Implied Volatility method.

The second part of this project is dedicated to modeling of implied volatility surface. We have studied two major classes of the volatility skews: the polynomial models (e.g., Cubic, Spline etc.) and stochastic models (the SABR model). Special attention was paid to the problems of the dynamics of the implied volatility surface in time. All of the models were tested on market data, which included different classes of underlying assets as well as different markets. We have suggested and applied different tests to compare implied volatility surface models. We have described the advantages and disadvantages of each method. We demonstrate that no single method exhibits superior accuracy in the analysis of every data set. Some methods perform better for certain underlying assets, while other methods are more suitable for the other.

5 Appendixes

5.1 *t* Location-Scale distribution

In this appendix we will discuss the *t* Location-Scale distribution, its likelihood function and the results of fitting the distribution to the returns of the datasets of Section 2.5.

The *t* Location-Scale distribution is a continuous distribution with the following density function:

$$f(x) = \frac{\Gamma\left(\frac{\nu+1}{2}\right)}{\sigma_{scale}\sqrt{\pi\nu}\Gamma\left(\frac{\nu}{2}\right)} \left[\frac{\nu + \left(\frac{x - \mu_{loc}}{\sigma_{scale}}\right)^2}{\nu} \right]^{-\frac{\nu+1}{2}}, \quad (5.1)$$

where

- Γ - gamma function,
- ν – shape parameter of the distribution,
- μ_{loc} - location parameter of the distribution,
- $\sigma_{scale} > 0$ – scale parameter.

If a random variable x has a *t* Location-Scale distribution, with parameters μ_{loc} , σ_{scale} and ν , then the following random variable

$$z = \frac{x - \mu_{loc}}{\sigma_{scale}},$$

will have a Student's *t* distribution with ν degrees of freedom.

The likelihood function of the *t* Location-Scale distribution is given by the following equation:

$$L(\theta, x) = f(x|\theta), \quad (5.2)$$

where

- $f(\cdot)$ - density function of a *t* Location-Scale distribution,
- $\theta = \{\nu, \mu_{loc}, \sigma_{scale}\}$ - parameters of the *t* Location-Scale distribution.

Values of parameters of *t* Location-Scale distribution for the testing datasets are given in Table 5-1.

Table 5-1. The values of parameters of the t Location-Scale distribution for different datasets

N	Data set	Location parameter: μ_{loc}	Scale parameter: σ_{scale}	Degrees of freedom: ν
1	Bonds	-0.00040	0.00868	3.03807
2	Commodities	0.00083	0.00976	2.03622
3	FX	0.00006	0.00508	4.33301
4	Indices	0.00033	0.00890	2.78986
5	Stocks	0.00003	0.01534	2.52348
6	Implied Volatility Indices	0.00033	0.00890	2.78986
7	Implied Volatility Stocks	0.00003	0.01534	2.52348

5.2 The Black-Scholes option pricing equation

In this appendix we introduce some definitions and give the Black-Scholes equation for option pricing.

An *European call option* is a financial contract that gives the holder the right but not the obligation to buy an underlying asset at a certain date (expiry date) for a certain price (exercise or strike price). *European put option*, unlike the call option, gives to its holder the right to sell an underlying asset at a certain date for a certain price.

In early 1970s, Fischer Black, Myron Scholes and Robert Merton made a major breakthrough into stock option pricing. This involved the development of what became known as the Black-Scholes model. In 1997 Myron Scholes and Robert Merton received a Nobel Prize in economics for their contribution in derivatives pricing. Let us introduce the assumptions and some important results of this model. The original assumptions are:

- The underlying stock price (S) is described by the following process:
$$\frac{dS}{S} = \mu dt + \sigma dW$$
 where μ is expected rate of return (the drift), σ is the constant volatility of the returns, W is a Brownian Motion.
- There are no transaction costs and all securities are perfectly divisible.
- There are no arbitrage opportunities.
- The risk free interest rate, r , is constant and the same for all maturities. Investors can freely borrow and lend money for the risk free interest rate.

The assumptions of the Black-Scholes model are rather strong and unrealistic. Extensions of the Black-Scholes model manage to overcome most of these restrictions. Still the assumption of constant volatility σ is one of the strongest.

Let us denote price of a European call option as C , P – European put and K strike price. Then C and P can be found by the following equations:

$$C = S_0 e^{-qT} N(d_1) - K e^{-rT} N(d_2), \quad (5.3)$$

$$P = K e^{-rT} N(-d_2) - S_0 e^{-qT} N(-d_1), \quad (5.4)$$

where,

$$d_1 = \frac{\ln(S_0/K) + (r - q + \sigma^2/2)T}{\sigma\sqrt{T}},$$

$$d_2 = d_1 - \sigma\sqrt{T},$$

$N(x)$ - is the cumulative probability distribution function for standard normal distribution,

S_0 - is the price of underlying asset at time $t = 0$,

q - is the expected dividend rate.

5.3 Cross-validation

In this appendix we discuss questions of dividing the data set into the subsets for validation of the results. Cross-validation is, probably, the simplest and the most effective method to estimate a prediction error on a data set (Hastie, Tibshirani, & Friedman, 2002). The main idea of this method is to divide the data set into several sub-parts, and to use some of them as a training set and others as a testing set. The training set is a part of all available data which is used to calibrate the models of interest, while the testing set is used to calculate the error.

The K –fold cross-validation could be described by the following algorithm. In the first step all available data is divided randomly into K equal parts (folds). So in each fold we have data which corresponds to different time periods. Then the 1st fold is used as a testing set and the remaining $K - 1$ folds are used as a training set. In the next step the 2nd fold is used as a testing set and the remaining folds (including the 1st one) are used as a training set. The algorithm is repeated on K folds. We will obtain K values of error; the resulting error is the average of these K values. We will take $K = 5$ for practical considerations.

The main advantage of the cross-validation technique is that the outliers and special observations will be equally distributed between the training and testing subsets. This is especially important for the problems of volatility estimation, because high values of volatility tend to cluster. If a certain model will be calibrated only on a low volatility period it will most probably fail to correctly estimate volatility on a high volatility testing set.

5.4 Results of the conditional volatility estimation

In this section we will present detailed results of the empirical tests of the volatility forecasting models discussed in Section 2.5. We will use the following notation for the characteristics of tests:

1. RMSE – Root Mean Square Error, see (2.11)
2. MHSE – Mean Heteroscedastic error , see (2.12)
3. $q_\alpha(Z)$ the quantile of the distribution of residuals Z of equation (2.6)
4. \widehat{ES}_q expected shortfall, see (2.31)
5. OES – Observed Expected Shortfall
6. CL – Confidence level

The results of the application of different volatility forecasting methods to different datasets are given in Tables 5-2 – 5-8.

Table 5-2. Performance of the volatility forecasting models on the Bond dataset.

N	Characteristic	Method					
		EWMA (λ^*)	GARCH (1,1)	ARCH(1)	EGARCH (1,1)	MB (RMSE)	MB (MHSE)
1	RMSE	0.943934	1.065585	1.095409	1.062676	0.900530	1.169438
2	MHSE	0.545672	1.542377	0.590675	2.442189	0.785110	0.430302
3	$q_\alpha(Z)$	4.320990	9.331957	4.726809	12.062141	5.144229	2.900205
4	\widehat{ES}_q	1.265371	1.396961	1.325958	1.735178	1.276882	1.241657
5	OES	1.276291	1.369750	1.305475	1.353536	1.294353	1.261523
6	CL	99.782501	99.752842	99.772615	99.782501	99.772615	99.752842

Table 5-3 . Performance of the volatility forecasting models on the Commodities dataset.

N	Characteristic	Method					
		EWMA (λ^*)	GARCH (1,1)	ARCH(1)	EGARCH (1,1)	MB (RMSE)	MB (MHSE)
1	RMSE	2.457830	2.651929	2.935664	2.631189	2.124783	3.826296
2	MHSE	1.039948	2.870677	0.854164	5.822701	2.249132	0.620857
3	$q_\alpha(Z)$	7.111687	13.154661	5.903431	19.707328	10.585036	3.579165
4	\widehat{ES}_q	1.702028	1.671963	1.633008	1.692778	1.552280	1.528889
5	OES	1.669192	1.691311	1.639904	1.709180	1.573506	1.536491
6	CL	99.782035	99.748012	99.707609	99.773530	99.715052	99.783099

Table 5-4. Performance of the volatility forecasting models on the FX dataset.

N	Characteristic	Method								
		EWMA (λ^*)	GARCH (1,1)	ARCH (1)	EGARCH (1,1)	GARCH (2,2)	ARCH (2)	EGARCH (2,2)	MB (RMSE)	MB (MHSE)
1	RMSE	0.483731	0.543138	0.522931	0.543906	0.524554	0.524554	0.529289	0.460114	0.601068
2	MHSE	0.505009	1.242811	0.555920	2.034977	0.863468	0.863468	1.086995	0.725140	0.415783
3	$q_\alpha(Z)$	4.191191	7.402903	4.316404	9.603921	5.717005	5.717005	6.632514	5.180113	2.992869
4	\widehat{ES}_q	1.326870	1.277245	1.359244	1.386234	1.239390	1.239390	1.364959	1.320630	1.328696
5	OES	1.060285	1.274581	1.309814	1.377977	1.245911	1.245911	1.362235	1.061135	1.297496
6	CL	99.845287	99.716351	99.699169	99.561619	99.690563	99.690563	99.630388	99.845291	99.836692

Table 5-5. Performance of the volatility forecasting methods on the Indices dataset.

N	Characteristic	Method								
		EWMA (λ^*)	GARCH (1,1)	ARCH (1)	EGARCH (1,1)	GARCH (2,2)	ARCH (2)	EGARCH (2,2)	MB (RMSE)	MB (MHSE)
1	RMSE	1.003088	1.083889	1.202130	1.066423	1.051140	1.051140	1.056888	0.926721	1.195491
2	MHSE	0.510880	1.099483	0.584689	1.928458	0.754898	0.754898	1.268886	0.849474	0.402964
3	$q_\alpha(Z)$	4.274652	6.785631	4.363502	8.017706	5.387333	5.387333	6.102579	5.770427	3.001282
4	\widehat{ES}_q	1.262509	1.329406	1.285756	1.575892	1.266997	1.266997	1.548677	1.310322	1.260705
5	OES	1.271655	1.332367	1.289575	1.579549	1.268335	1.268335	1.510626	1.314619	1.265185
6	CL	99.796638	99.730424	99.661849	99.628742	99.735156	99.735156	99.678405	99.827378	99.839201

Table 5-6. Performance of the volatility forecasting methods on the Stocks dataset.

N	Characteristic	Method					
		EWMA (λ^*)	GARCH (1,1)	ARCH(1)	EGARCH (1,1)	MB (RMSE)	MB (MHSE)
1	RMSE	2.006486	2.197144	2.424559	2.186051	1.865170	2.583198
2	MHSE	0.604923	1.553748	0.625334	14.168314	0.931479	0.470888
3	$q_\alpha(Z)$	5.094329	8.774326	5.053610	11.026071	6.574708	3.423447
4	\widehat{ES}_q	1.436950	1.432639	1.366123	2.906567	1.450504	1.442305
5	OES	1.436849	1.434995	1.367075	2.923439	1.449347	1.449301
6	CL	99.810792	99.732023	99.796023	99.704454	99.810463	99.822607

Table 5-7. Performance of the volatility models and implied volatility for the Indices data subset.

N	Characteristic	Method									
		EWMA (λ^*)	GARCH (1,1)	ARCH (1)	EGARCH (1,1)	GARCH (2,2)	ARCH (2)	EGARCH (2,2)	MB (RMSE)	MB (MHSE)	IV ¹
1	RMSE	1.090791	1.171080	1.388507	1.138602	1.137115	1.137115	1.127102	0.991736	1.286520	1.062791
2	MHSE	0.546538	1.013394	0.600877	1.791266	0.727686	0.727686	1.033894	1.062797	0.420414	0.511294
3	$q_\alpha(Z)$	4.624267	7.299857	4.982415	9.831520	5.750332	5.750332	7.443971	7.290430	3.038510	4.257364
4	\widehat{ES}_q	1.222764	1.133752	1.191009	1.561176	1.130966	1.130966	1.280882	1.560017	1.167447	1.189892
5	OES	0.468067	0.907141	0.702527	1.544002	0.674703	0.674703	0.772661	0.477010	0.467509	0.952244
6	CL	99.8344	99.7681	99.8012	99.7350	99.7847	99.7847	99.8012	99.8675	99.8675	99.7846

Table 5-8. Performance of the volatility models and implied volatility for the Stocks data subset.

N	Characteristic	Method									
		EWMA (λ^*)	GARCH (1,1)	ARCH (1)	EGARCH (1,1)	GARCH (2,2)	ARCH (2)	EGARCH (2,2)	MB (RMSE)	MB (MHSE)	IV ²
1	RMSE	3.636001	3.642840	4.035042	3.408410	3.649005	3.649005	3.445134	3.237717	5.684578	3.201078
2	MHSE	1.296614	1.726116	0.881681	2.474553	1.530718	1.530718	1.608248	1.908309	0.806629	1.223559
3	$q_\alpha(Z)$	5.808708	7.311894	5.489278	8.973477	6.704227	6.704227	6.956462	7.383313	2.833592	5.918723
4	\widehat{ES}_q	2.616192	1.905565	1.690664	2.016285	2.101178	2.101178	1.883677	2.338671	1.934981	2.181829
5	OES	1.540882	1.161122	1.360862	1.075147	0.973995	0.973995	1.134165	1.179503	1.857380	1.294764
6	CL	99.8438	99.7321	99.7098	99.7098	99.7545	99.7545	99.6875	99.8438	99.7321	99.7545

¹ Implied volatility is not included in Model Blending procedures.

5.5 Results of implied volatility surface modeling

In this appendix we present the results of the modeling volatility skews for each of the underlying assets. The errors for different types of tests and underlying are presented in Tables 5-9 – 5-17.

Table 5-9. Performance of different volatility surface models on options of the AEX index.

N	Model	Quality of fit	Static models		Dynamic models	
			1 day forecast	5 days forecast	1 day forecast	5 days forecast
1	Cubic	1.28626	3.21429	6.37292	3.99260	13.4539
2	Spline	1.05448	2.61850	5.39306	3.78533	9.41520
3	SABR	3.88212	5.62535	8.00410	5.62627	8.37488
4	PSABR	1.64033	3.48644	6.29141	6.05896	8.83062
5	Sticky Strike	N/A	1.97364	4.36168	N/A	N/A

Table 5-10. Performance of different volatility surface models on options of the German stock index DAX.

N	Model	Quality of fit	Static models		Dynamic models	
			1 day forecast	5 days forecast	1 day forecast	5 days forecast
1	Cubic	1.34813	3.25822	5.99786	3.48532	7.19360
2	Spline	0.84351	1.96317	4.06831	2.01622	4.57051
3	SABR	4.00186	5.08159	6.37802	5.08352	6.70261
4	PSABR	1.89345	2.83703	4.86227	4.74022	6.49834
5	Sticky Strike	N/A	1.21387	2.95181	N/A	N/A

Table 5-11. Performance of different volatility surface models on options of the European stock index EURSTOXX.

N	Model	Quality of fit	Static models		Dynamic models	
			1 day forecast	5 days forecast	1 day forecast	5 days forecast
1	Cubic	0.33500	1.65018	4.55425	1.82739	5.44496
2	Spline	0.31123	1.66325	4.56096	2.16492	5.96404
3	SABR	4.76126	5.18839	8.44978	5.19007	7.33259
4	PSABR	2.89203	3.40392	6.20050	3.64290	5.71112
5	Sticky Strike	N/A	0.87453	2.38508	N/A	N/A

Table 5-12. Performance of different volatility surface models on options of the FTSE 100 – the UK equity index.

N	Model	Quality of fit	Static models		Dynamic models	
			1 day forecast	5 days forecast	1 day forecast	5 days forecast
1	Cubic	0.33445	1.65999	4.58970	1.84163	5.50039
2	Spline	0.31083	1.67336	4.59633	2.17110	5.94110
3	SABR	4.26895	4.56957	8.48371	4.57191	6.89312
4	PSABR	2.86000	3.39127	6.17133	3.58511	5.79384
5	Sticky Strike	N/A	0.87875	2.39056	N/A	N/A

Table 5-13. Performance of different volatility surface models on options of the XJO index – the Australian equity index.

XJO						
N	Model	Quality of fit	Static models		Dynamic models	
			1 day forecast	5 days forecast	1 day forecast	5 days forecast
1	Cubic	2.97195	11.01804	11.60088	9.99524	12.51182
2	Spline	2.87676	9.92295	10.63055	7.91253	8.84867
3	SABR	9.13593	13.43551	14.81607	13.42370	14.45272
4	PSABR	4.13998	9.56873	11.04536	11.20640	12.45780
5	Sticky Strike	N/A	6.45673	7.91869	N/A	N/A

Table 5-14. Performance of different volatility surface models on implied volatility of the Deutsche Bank shares.

DBK						
N	Model	Quality of fit	Static models		Dynamic models	
			1 day forecast	5 days forecast	1 day forecast	5 days forecast
1	Cubic	7.74958	17.13035	36.88834	50.52760	136.69437
2	Spline	6.96815	14.86619	33.51260	28.45704	76.44249
3	SABR	16.68262	26.54118	55.61741	26.56520	46.13593
4	PSABR	8.78661	17.59187	35.20103	21.83207	35.48460
5	Sticky Strike	N/A	13.59281	27.14709	N/A	N/A

Table 5-15. Performance of different volatility surface models on implied volatility of the Volkswagen shares.

VOW						
N	Model	Quality of fit	Static models		Dynamic models	
			1 day forecast	5 days forecast	1 day forecast	5 days forecast
1	Cubic	3.06962	9.36626	30.12601	15.53513	61.48214
2	Spline	3.09983	9.36353	31.45669	10.79304	44.93108
3	SABR	30.62417	44.06871	75.15178	44.08075	72.31071
4	PSABR	6.16194	13.91551	37.20842	14.84501	34.54480
5	Sticky Strike	N/A	7.33491	26.04306	N/A	N/A

Table 5-16. Performance of different volatility surface models on implied volatility of the Allianz shares.

ALV						
N	Model	Quality of fit	Static models		Dynamic models	
			1 day forecast	5 days forecast	1 day forecast	5 days forecast
1	Cubic	6.31670	12.95525	25.57154	14.91010	31.68649
2	Spline	6.05479	11.29525	21.63830	13.15205	33.84061
3	SABR	11.63363	16.75927	34.90202	16.78393	27.27270
4	PSABR	6.87484	11.46044	23.75863	13.58053	24.16736
5	Sticky Strike	N/A	7.91247	15.41104	N/A	N/A

Table 5-17. Performance of different volatility surface models on implied volatility of the Adidas shares.

ADS						
N	Model	Quality of fit	Static models		Dynamic models	
			1 day forecast	5 days forecast	1 day forecast	5 days forecast
1	Cubic	0.71324	2.54451	7.69154	2.65212	9.37217
2	Spline	0.70272	2.25693	6.82459	2.76832	9.73861
3	SABR	2.92377	4.72371	9.83645	4.72329	9.33579
4	PSABR	1.42065	3.17030	7.97566	3.77238	8.49589
5	Sticky Strike	N/A	1.83600	5.51145	N/A	N/A

6 Bibliography

Black, F., & Scholes, M. (1973). The Pricing of Options and Corporate Liabilities. *Journal of Political Economy* , 637–654.

Engle, R. (1982). Autoregressive conditional heteroscedasticity with estimates of the variance of United Kingdom inflation. *Econometrica* , 987–1007.

Hafner, R., & Wallmeier, M. (2000). The Dynamics of DAX Implied Volatilities.

Hagan, P., Kumar, D., Lesniewski, A., & Woodward, D. (2002). Managing Smile Risk. *Wilmott Magazine* , 84-108.

Hastie, T., Tibshirani, R., & Friedman, J. (2002). *The Elements of Statistical Learning*. Springer Series in Statistics.

Hull, J. C. (2002). *Options, futures and other derivatives*. New Jersey: Prentice Hall.

J. Knight, S. S. (2007). *Forecasting volatility in the financial markets*. London: Elsevier.

J.P. Morgan/Reuters. (1996). *RiskMetrics - Technical Document*. New York: J.P. Morgan/Reuters.

McNeil, A. (1999). Extreme Value Theory for Risk Managers.

McNeil, F. &. (2005). *Quantitative Risk Management: Concepts, Techniques, and Tools*. Princeton University Press.

Nelson, D. (1991). Conditional Heteroskedasticity in Asset Returns: A New Approach. *Econometrica* , 347–370.

Poon, S. H. (2005). *A practical guide to forecasting financial market volatility*. Chichester: John Wiley and Sons, Ltd.

Poon, S. (2008). SABR.

Shlens, J. (2005). *A Tutorial on Principal Component Analysis*.

Tsai, H. (2006). *A Note on Inequality Constraints in the GARCH model*. Iowa: The University of Iowa.

Tsay, R. S. (2005). *Analysis of Financial Time Series*. Wiley-Interscience; 2nd edition .

West, G. (2005). Calibration of the SABR Model in Illiquid Markets. *Applied Mathematical Finance* , 371-385.

Wilmott, P., Howison, S., & Dewynne, J. (1995). *The Mathematics of Financial Derivatives*. Cambridge: Cambridge University Press.

Witten, H. (2005). *Data Mining: practical machine learning tools and techniques*. Elsevier.